Resonant Leptogenesis and Collider Signals from Discrete Flavor and CP Symmetries

Garv Chauhan Washington University in St. Louis, USA

PPC 2021: XIV International Workshop on Interconnections between Particle Physics and Cosmology May 20, 2021

Work based on GC and Bhupal Dev, arXiv:2106.abcde





[Fukugita, Yanagida (Phys. Lett. B '86)]

- Central idea : Leptonic asymmetry in early Universe is converted to baryonic asymmetry through B-L conserving EW sphaleron interactions.
- Add SM-singlet heavy Majorana neutrinos.

$$\mathcal{L}_l = Y_l \bar{L}_l H l_R + Y_D \bar{L}_l \tilde{H} N + \frac{1}{2} \bar{N}^c M_R N + h.c.$$

- Satisfies all 3 Sakharov conditions.
 - CP violation in the leptonic sector (through complex Y_D and/or U_{PMNS} phases)
 - L violation due to the Majorana nature of the heavy RH neutrinos
 - Departure from thermal equilibrium when $\Gamma_N \leq H$
- It can connect neutrino mass mechanism and matter-antimatter asymmetry.



- TeV scale leptogenesis, no dependence on initial conditions.
- If $\Delta m_N \sim \Gamma_N \ll m_N$, the self energy contribution (ε -type) to the CP asymmetry becomes dominant and large (even order 1).
- The ε -type CP asymmetry,

$$\varepsilon_{N_i} = \frac{\mathrm{Im}(h^{\nu\dagger}h^{\nu})_{ij}^2}{(h^{\nu\dagger}h^{\nu})_{ii}(h^{\nu\dagger}h^{\nu})_{jj}} \frac{(M_{N_i}^2 - M_{N_j}^2)M_{N_i}\Gamma_{N_j}}{(M_{N_i}^2 - M_{N_j}^2)^2 + M_{N_i}^2\Gamma_{N_j}^2}$$

• Order 1 CP aymmetries are possible when, [Pilaftsis '97; Pilaftsis, Underwood '03]

$$M_{N_2} - M_{N_1} \sim \frac{1}{2} \Gamma_{N_{1,2}}$$
$$\frac{\ln(h^{\nu^{\dagger}} h^{\nu})_{ij}^2}{(h^{\nu^{\dagger}} h^{\nu})_{ii} (h^{\nu^{\dagger}} h^{\nu})_{jj}} \sim 1$$

• This helps lower the heavy neutrino scale M_N , which can be as low as EW scale. [Pilaftsis, Underwood '05; Deppisch, Pilaftsis '10; Dev, Millington, Pilaftsis, Teresi '14]

- High energy neutrino parameters are free parameters in the leptogenesis mechanism.
- We will look at the idea of residual flavor and CP symmetries that determine lepton mixing angles, low- and high energy CP phases with only one free parameter.
- We conjecture the existence of a finite, discrete flavor symmetry G_f at a high-energy scale
- At low energies, G_f is broken to G_l in the charged lepton sector and to G_ν in the neutrino sector.
- G_l determines U_l and G_{ν} determines U_{ν} . This leads to the PMNS matrix

$$U_{PMNS} = U_l^{\dagger} U_{\nu}$$

- For G_f , we use a group of the form $\Delta(6n^2)$ with n even , $3 \nmid n$ and $4 \nmid n$..
- Residual symmetries : $G_l = Z_3$, $G_\nu = Z_2 \times CP$
- given X (CP transformation) and Z (generator of Z_2 in 3)

 $Z^{\dagger}(\mathbf{3}) \, Y_D \, Z(\mathbf{3}') = Y_D \quad \text{and} \quad X^{\star}(\mathbf{3}) \, Y_D \, X(\mathbf{3}') = Y_D^{\star} \; .$

Consistency condition $:X(\mathbf{r}) Z(\mathbf{r}) = Z(\mathbf{r})^* X(\mathbf{r})$

 \bullet Changing to a different basis by the unitary matrix Ω that fulfills

 $\Omega^{\dagger} Z \Omega = \mathsf{diag}((-1)^{z_1}, (-1)^{z_2}, (-1)^{z_3}) \ z_i = 0, 1$

it follows then $X = \Omega \Omega^T$ and $\Omega^T Y_D \Omega$ real.

• $\Omega^T Y_D \Omega$ can be diagonalized by two rotation matrices from the left and right, respectively

$$\Omega(s)(\mathbf{3})^{\dagger} Y_D \Omega(s)(\mathbf{3}') = R_{ij}(\theta_L) \begin{pmatrix} y_1 & 0 & 0\\ 0 & y_2 & 0\\ 0 & 0 & y_3 \end{pmatrix} R_{kl}(-\theta_R).$$

• $U_{\text{PMNS}} = \Omega(\mathbf{3}) R_{ij}(\theta_L) K_{\nu}, \qquad K_{\nu} = \begin{pmatrix} 1 & 0 & 0\\ 0 & i^{k_1} & 0\\ 0 & 0 & i^{k_2} \end{pmatrix} k_i = 0, 1, 2, 3$

ullet The light neutrino mass matrix m_{ν} follows from the type-I seesaw mechanism

$$m_{\nu} = m_D M_R^{-1} m_D^T \,.$$

Our chosen case : $\Delta(6n^2)$

٠

 $\mathbf{m}_{\nu}: \left\{ \begin{array}{l} \frac{1}{M_{N}} \begin{pmatrix} y_{1}^{2}\cos 2\theta_{R} & 0 & y_{1}y_{3}\sin 2\theta_{R} \\ 0 & y_{2}^{2} & 0 \\ y_{1}y_{3}\sin 2\theta_{R} & 0 & -y_{3}^{2}\cos 2\theta_{R} \end{pmatrix} \right. \quad \mathbf{s \ even} \\ \left. \begin{array}{l} \frac{1}{M_{N}} \begin{pmatrix} -y_{1}^{2}\cos 2\theta_{R} & 0 & -y_{1}y_{3}\sin 2\theta_{R} \\ 0 & y_{2}^{2} & 0 \\ -y_{1}y_{3}\sin 2\theta_{R} & 0 & y_{3}^{2}\cos 2\theta_{R} \end{array} \right) \mathbf{s \ odd} \end{array} \right.$

• For $y_1 = 0 (y_3 = 0)$, we get strong normal (inverted) ordering, with $m_{\text{lightest}} = 0$.

$$NO: \quad y_{1} = 0, \quad y_{2} = \pm \frac{\sqrt{M_{N}\sqrt{\Delta m_{sol}^{2}}}}{v}, \quad y_{3} = \pm \frac{\sqrt{M_{N}\frac{\sqrt{\Delta m_{atm}^{2}}}{|\cos 2\theta_{R}|}}}{v}$$
$$IO: \quad y_{3} = 0, \quad y_{2} = \pm \frac{\sqrt{M_{N}\sqrt{\Delta m_{atm}^{2}}}}{v}, \quad y_{1} = \pm \frac{\sqrt{M_{N}\frac{\sqrt{|\Delta m_{atm}^{2}| - \Delta m_{sol}^{2}}}{|\cos 2\theta_{R}|}}}{v}$$

• Only free parameters : M_N and θ_R

Collider Signal

 $\bullet\,$ In our scenario, $y_i \lesssim 10^{-6}$ supresses the Drell Yan production

$$pp \to W^{(*)} \to N_i l_{\alpha}$$

- We need to go beyond the minimal type-I seesaw to realize a sizable LNV signal.
- This scenario can also be embedded in SM with extended gauge symmetry
- We consider minimal $U(1)_{B-L}$ extension for enhanced production of N_i at colliders.



Collider Signal - Branching Ratio



For $M_{Z'} = 4$ TeV and s = 2, n = 26



• The decay widths Γ_i of the RH neutrinos N_i are given at the tree level by

$$\Gamma_i \approx \frac{(\hat{Y}_D^{\dagger} \, \hat{Y}_D)_{ii}}{8 \, \pi} M_i = \frac{(\hat{m}_D^{\dagger} \, \hat{m}_D)_{ii}}{8 \, \pi \, v^2} M_i$$

• The expressions for decay widths of the 3 heavy RH neutrinos :

$$\begin{split} \Gamma_1 &\approx \quad \frac{M}{24\,\pi}\,\left(2\,y_1^2\,\cos^2\theta_R + y_2^2 + 2\,y_3^2\,\sin^2\theta_R\right)\\ \Gamma_2 &\approx \quad \frac{M}{24\,\pi}\,\left(y_1^2\,\cos^2\theta_R + 2\,y_2^2 + y_3^2\,\sin^2\theta_R\right)\,,\\ \Gamma_3 &\approx \quad \frac{M}{8\,\pi}\,\left(y_1^2\,\sin^2\theta_R + y_3^2\,\cos^2\theta_R\right)\,. \end{split}$$

- If $\theta_R \approx \pi/2$, $3\pi/2$ (for strong NO) or $\theta_R \approx 0$, π (for strong IO), Γ_3 tends to zero. (termed Enhanced Residual Symmetry points)
- Near points of ERS , N_3 can have a very long lifetime $\rightarrow N_3$ may be detected in long-lived particle searches such as in MATHUSLA detector.





 $\theta_R \approx 0, \pi$ (ERS points)

- At leading order, we have three degenerate RH neutrinos.
- Higher-order corrections can break the residual symmetries :

 $M_1 = M_N (1 + 2 \kappa)$ and $M_2 = M_3 = M_N (1 - \kappa)$.

• CP asymmetries in the decays of N_i are given by :

$$\epsilon_{i\alpha} \sim \sum_{j} \operatorname{Im}(\hat{Y}_{D,\alpha i}^{*} Y_{D,\alpha i}) \operatorname{Re}(\hat{Y}_{D}^{\dagger} Y_{D})_{ij} F_{ij}$$

- F_{ij} are related to the regulator in ReL and are proportional to the mass splitting of N_i .
- We find

$$\varepsilon_{1\alpha} \sim \frac{y_2 y_3}{9} (-2y_2^2 + y_3^2 (1 - \cos 2\theta_R)) \sin 3\phi_s \sin 2\theta_R \sin \theta_{L,\alpha} F_{12} \text{ (NO)}$$

$$\varepsilon_{1\alpha} \sim \frac{y_1 y_2}{9} (-2y_2^2 + y_1^2 (1 + \cos 2\theta_R)) \sin 3\phi_s \sin 2\theta_R \sin \theta_{L,\alpha} F_{12} \text{ (IO)}$$

with $\theta_{L,\alpha}=\theta_L+\rho_\alpha 4\pi/3$ and $\rho_e=0,\rho_\mu=1,\rho_\tau=-1$



$$g_{B-L} = 0.1, s = 2, n = 26$$

and $\theta_R = \frac{\pi}{2} (0)$ for strong NO (IO)

 $\frac{\varepsilon_{NO}}{\varepsilon_{IO}} = \left(\frac{\Delta m^2_{\rm atm}}{\Delta m^2_{\rm sol}}\right)^{3/4}$



For $g_{B-L} = 0.1$, $M_{Z'} \gtrsim 4.12 \text{ TeV}$ For $g_{B-L} = 1$, $M_{Z'} \gtrsim 7 \text{ TeV}$



all cross sections in ab

For $g_{B-L} = 1$, $M_{Z'} \gtrsim 7 \text{ TeV}$ Enhancement from $g_{B-L}^4 \rightarrow 10^4$

- Neutrinoless double beta $(0\nu\beta\beta)$ decay is one of the most important theorised LNV process to discern the Majorana nature of the neutrinos.
- The predictions for this yet unobserved process depends explicitly on the Majorana phases α and β .
- A nuclear isotope decaying through $0\nu\beta\beta$ decay would exhibit a half-life $T_{1/2}^{0\nu\beta\beta}$ of

$$\Gamma^{0\nu\beta\beta} = \frac{1}{T_{1/2}^{0\nu\beta\beta}} = G^{0\nu} |M^{0\nu}|^2 \frac{m_{ee}}{m_e}$$

$$m_{ee} = \left| U_{\rm PMNS,11}^2 \, m_1 + U_{\rm PMNS,12}^2 \, m_2 + U_{\rm PMNS,13}^2 \, m_3 \right|$$
$$m_{ee} = \left| \cos^2 \theta_{12} \, \cos^2 \theta_{13} \, m_1 + \sin^2 \theta_{12} \, \cos^2 \theta_{13} \, e^{i\alpha} \, m_2 + \sin^2 \theta_{13} \, e^{i\beta} \, m_3 \right| \,.$$

 $\bullet\,$ In our example case, the light neutrino contribution to $0\nu\beta\beta$ is restricted to :

$$\mathbf{m}_{\beta\beta}: \frac{1}{3} \begin{cases} \left| \sqrt{\Delta m_{\mathsf{sol}}^2} + 2(-1)^{s+k+1} \sin^2 \theta_L e^{6i\phi_s} \sqrt{\Delta m_{\mathsf{atm}}^2} \right| & \text{(NO)} \\ \\ \left| 1 + 2(-1)^{s+k+1} \cos^2 \theta_L e^{6i\phi_s} \right| \sqrt{|\Delta m_{\mathsf{atm}}^2|} & \text{(IO)} \end{cases}$$

$0\nu\beta\beta$ results



- Leptogenesis is an attractive mechanism to explain the BAU.
- Resonant Leptogenesis leads to order 1 *CP* asymmetry and reduces the energy scale of BAU production to TeV scale.
- The high-energy *CP* violating physics is disconnected from low-energy neutrino data, can be connected through role of residual flavor and CP symmetries.
- We have presented a type-I seesaw scenario with a flavour and CP symmetry as well as three RH neutrinos with almost degenerate masses in the few hundred GeV to TeV range.
- Requiring η_B to be generated via resonant leptogenesis constrains the prospects for detecting RH neutrinos at colliders
- Tight predictions for future neutrinoless double beta decay experiments can fully probe our scenario and thus provide complementary information

Supplementary Material

Leptogenesis - 3 basic steps

• Generation of L asymmetry in heavy Majorana neutrino N decay :



• Partial washout of the asymmetry due to inverse decay and scatterings with $\Delta L \neq 0$:



• Conversion of the leftover L asymmetry to B asymmetry at $T > T_{sph}$:



Our chosen case : $\Delta(6n^2)$

- For G_f , we use a group of the form $\Delta(6n^2)$ with n even , $3 \nmid n$ and $4 \nmid n$..
- Residual symmetries : $G_l = Z_3$, $G_\nu = Z_2 \times CP$
- $\Delta(6n^2) = (Z_n \times Z_n) \rtimes S_3$ $a^3 = e, \ c^n = e, \ d^n = e, \ cd = dc, \ a \, c \, a^{-1} = c^{-1} d^{-1}, \ a \, d \, a^{-1} = c$ $b^2 = e, \ (a \, b)^2 = e, \ b \, c \, b^{-1} = d^{-1}, \ b \, d \, b^{-1} = c^{-1}.$

• For case in consideration : $Z = c^{n/2}$ and $X = a \, b \, c^s \, d^{2s}$ with s = 0, 1, ..., n-1

• As M_R leaves G_f and CP invariant, its form is simply

$$M_R = M_N \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array} \right) \tag{1}$$

- Dirac CP phase is trivial $\delta = 0$.
- For $m_{\text{lightest}} = 0$, only one Majorana phase α , which depends on the chosen CP transformation:

$$\sin \alpha = (-1)^{k+r+s} \sin 6\phi_s$$
 and $\cos \alpha = (-1)^{k+r+s+1} \cos 6\phi_s$ with $\phi_s = \frac{\pi s}{n}$

where k = 0(k = 1) for $\cos 2\theta_R > 0(\cos 2\theta_R < 0)$ and r = 0(r = 1) for NO(IO).